

Undirected Graphs of Entanglement 3

Walid Belkhir

walid.belkhir@lif.univ-mrs.fr

Laboratoire d'Informatique Fondamentale de Marseille.

Université de Provence.

Abstract

Entanglement is a complexity measure of digraphs that originates in fixed-point logics. Its combinatorial purpose is to measure the nested depth of cycles in digraphs. We address the problem of characterizing the structure of graphs of entanglement at most k . Only partial results are known so far: digraphs for $k = 1$, and undirected graphs for $k = 2$. In this paper we investigate the structure of undirected graphs for $k = 3$. Our main tool is the so-called *Tutte's decomposition* of 2-connected graphs into cycles and 3-connected components into a tree-like fashion. We shall give necessary conditions on Tutte's tree to be a tree decomposition of a 2-connected graph of entanglement 3.

Key words. Entanglement, tree decomposition, Tutte's decomposition, connectivity.

1 Introduction

Entanglement is a complexity measure of digraphs introduced in [BG05] to analyze the descriptive complexity of the propositional modal μ -calculus [Koz83]. This measure has shown its use in solving the variable hierarchy problem¹ for the modal μ -calculus [BGL07] and for the lattice μ -calculus [BS08]. However, little is known about the graph theoretic and algorithmic properties of entanglement, such that the structural properties of graphs of bounded entanglement. The structure of digraphs of entanglement ≤ 1 is known [BG05]. Using this result, it was argued that deciding whether a digraph has entanglement ≤ 1 is a problem in NLOGSPACE. The graphs² of entanglement ≤ 2 have been characterized in [BS07]. Using this characterization, a linear time algorithm recognizing those graphs has been devised.

In this paper we study the structure of graphs of entanglement 3. Our approach consists in studying entanglement w.r.t. the notions of connectivity and

¹This problem asks whether the expressive power of a given fixed-point logic increases with the number of bound variables.

²Throughout this paper, an undirected graph is called simply a graph; a directed graph is called a digraph.

cyclicity. The latter are among the basic properties of graphs: a graph may be decomposed into maximal articulation point-free components in a tree like fashion. An articulation point-free component itself admits a canonical decomposition in terms of cycles and 3-connected components in a tree like structure, this decomposition was given by Tutte [Tut66]. Continuing this process, Høhberg gave a decomposition of graphs of higher connectivity [Hoh92], this is a sort of generalization of Tutte's decomposition. However, Høhberg's decomposition is no longer canonical. Another decomposition of digraphs in terms of directed cycles, known as *ear* decomposition [BJG01, §7], has been useful for studying the connectivity of directed multigraphs. Such decompositions are important tools in graph theory and they are often used in inductive proofs and constructions. For instance see [RSS94] where Tutte's decomposition is used.

In [BS07] it was pointed out that the canonical decomposition of graphs into 2-connected components is well adapted in characterizing the structure of graphs of entanglement ≤ 2 . With a similar approach we are asking whether Tutte's decomposition would be of use to characterize the structure of graphs of entanglement 3. Roughly speaking Tutte's decomposition reduces the structure of a given graph into that of its components. So what can we say about the components themselves if the given graph has entanglement 3? We shall find necessary conditions on Tutte's tree to be a tree decomposition of a 2-connected graph of entanglement 3: we shall (i) characterize the structure of 3-connected components of G , (ii) impose necessary conditions on the manner by which those components are glued together, and (iii) find an upper bound of the diameter of Tutte's tree.

We suggest a natural path for generalizing the algebraic approach used to construct the graphs of entanglement 2 [BS07] to the construction of the graphs of entanglement 3. We mean by the algebraic approach the manner by which a class of graphs is build up out of small pieces of graphs using an appropriate set of gluing operations. This generalization interacts with Tutte's decomposition in a deep and not obvious way. Finally, ear decomposition would be useful in dealing with the entanglement within the direct setting. Furthermore, Høhberg's decomposition suggests a hopeful path for the characterization of graphs of higher entanglement.

2 Preliminaries

A directed graph $G = (V_G, E_G)$ is a set of vertices V_G and a binary relation $E_G \subseteq V_G \times V_G$. All the graphs in this paper are finite and undirected until we say otherwise. They are also simple i.e. without multi-edges, but during the decomposition process³ some multi-edges may appear. If $X \subseteq V_G$, then we shall write $G[X]$ for the subgraph of G induced by X .

³Indeed we mean Tutte's decomposition.

2.1 Cyclicity

A *feedback vertex set* [FPR99] of a digraph G is a subset $X \subseteq V_G$ that meets all directed cycles of G , i.e. the digraph $G \setminus X$ is acyclic. The *cyclicity* of G , denoted $\mathcal{C}(G)$, is the cardinality of the *minimum* feedback vertex set.

When we deal with the cyclicity of a graph G we consider the *directed* cycles of the *symmetric directed* graph G . In other words, every (undirected) edge v_1v_2 of G may be viewed as the set of (directed) edges $\{(v_1, v_2), (v_2, v_1)\}$, and therefore $v_1v_2v_1$ is a directed cycle of length 2. To make the notion of the cyclicity of a graph easier and more intuitive, it is convenient to see the feedback vertex set of a graph as an *edge cover set*, i.e. a set $X \subseteq V_G$ is an edge cover of G if for each edge $v_1v_2 \in E_G$ we have that $v_1 \in X$ or $v_2 \in X$.

Lemma 2.1. *Let G be an undirected graph, and let $X \subseteq V_G$. Then, X is a feedback vertex set of G if and only if X is an edge cover set of G .*

Proof.

$\Rightarrow$

Let $X \subseteq V_G$ be a feedback vertex set of G . Therefore, for every cycle of the form $v_1v_2v_1$ we have that $v_1 \in X$ or $v_2 \in X$. Since the cycle $v_1v_2v_1$ is just the edge v_1v_2 of the undirected graph G , then X is an edge cover set of G .

$\Leftarrow$

Let $X \subseteq V_G$ be an edge cover set of G and let $C_n = v_1v_2 \dots v_nv_1$ be a cycle of G . Then, for each edge v_iv_{i+1} of C_n we have $v_i \in X$ or $v_{i+1} \in X$, because X is an edge cover. Therefore X meets C_n , hence X is a feedback vertex set of G . $\square$

It follows by the previous Lemma that the cyclicity of an undirected graph G is the cardinality of the minimum edge cover set of G .

2.2 Connectivity

Given two sets A, B of vertices, we call $\pi = v_1 \dots v_n$ an *A - B path* if $V_\pi \cap A = \{v_1\}$ and $V_\pi \cap B = \{v_n\}$. Two or more paths are *independent* if none of them contains an internal vertex of the other. If a, b are two vertices then we shall write " *a - b path*" instead of " $\{a\}$ - $\{b\}$ path". In this case, two *a - b paths* are independent if and only if a and b are their only common vertices.

Definition 2.2. A graph is *k -connected* [Die05, §3] if one needs to remove at least k -vertices to disconnect it. The *connectivity* of a graph G is the maximum k such that G is k -connected.

The following Theorem, due to Menger [Die05, §3.3], provides a useful equivalent definition of k -connectivity.

Theorem 2.3. *A graph is k -connected if and only if it contains k independent paths between any two vertices.*

A vertex whose removal disconnects the graph is called *articulation point*. A maximal connected subgraph without articulation points is called a *biconnected component*. By their maximality, the biconnected components of G overlap in at most one vertex, which is an articulation point of G . Hence, every edge of G belongs to a unique biconnected component. An isolated vertex is considered trivially as a biconnected component. Let $\mathcal{A}$ be the set of articulation points of G , and $\mathcal{B}$ the set of its 2-connected components. Then we obtain a bipartite graph $T_1(G)$ whose vertices are $\mathcal{A} \cup \mathcal{B}$ and whose edges are aB whenever $a \in B$. It is one of the elementary results to show that if G is connected then $T_1(G)$ is a tree, see Proposition 3.1.2 of [Die05].

2.3 Separations, hinges

In this subsection we introduce the definition of separations and hinges. The material presented in this here can be found in [Tut66], whereas we shall adopt the notation and the terminology of [DSS95] and [Ric04].

A *bond* is a graph consisting of just two vertices and at least one edge. A *k-bond* is a bond of k edges. Observe that, if $k \geq 2$, then a k -bond is a multi-graph. If $A, B \subseteq V_G$ and $S \subset V_G$ are such that every A - B path in G contains a vertex of S , then we say that S separates the sets A and B in G . Observe that this implies that $A \cap B \subseteq S$. We shall say that S separates G if $G \setminus S$ is disconnected, that is, if S separates G into some vertices which are not in S . A separating set of vertices is called a *separator*. A pair (A, B) is a *separation* of G if $A \cup B = V_G$ and G has no edge between $A \setminus B$ and $B \setminus A$. Clearly, this is equivalent to saying that $A \cap B$ separates A from B . We say that a separation (H, K) is a *k-separation* if $|V_{H \cap K}| = k$. A subgraph K of G is an *H-bridge* if K is obtained from a component $\mathcal{C}$ of $G \setminus V_H$ by adding to $\mathcal{C}$ all the edges of G which have at least one end in $\mathcal{C}$. From the above definition of connectivity follows a standard result of graph theory [Tut66]:

Lemma 2.4. *A graph is k -connected if there is no m -separation, for all $m = 0, \dots, k - 1$.*

From the definition of connectivity, it follows that the connectivity of the following graphs is infinite: the graph of just one vertex, the k -bonds, for $k \geq 1$, and the 3-clique, see Figure 2.1.



Figure 2.1: Graphs with infinite connectivity.

Convention 2.5. *For technical reasons, we shall consider the connectivity of the k -cliques, for $k \geq 3$, equals $k - 1$.*

If (A, B) is a 2-separation of G , then we say that $A \cap B$ is a *hinge* of G iff at least one of $G[A]$ and $G[B]$ is 2-connected.

2.4 Graph minors

A graph G is *minor* of a graph H if G can be obtained from H by successive application of the following operations on it:

- (a) delete an edge,
- (a) contract an edge, and
- (a) delete an isolated vertex.

A class $\mathcal{C}$ of graphs is minor closed if it is closed under taking minors, that is whenever $H \in \mathcal{C}$ then for each minor G of H we have that $G \in \mathcal{C}$.

2.5 Tree decomposition

We reproduce here the definition of the notion of *tree decomposition* as given by Robertson and Seymour [RS86].

Definition 2.6. Let G be a graph, T be a tree, and let $X = (V_t)_{t \in T}$ be a family of subsets of V_G . We say that the pair (T, X) is a *tree decomposition* of G if the following conditions hold:

- (T-1) $V_G = \bigcup_{t \in T} V_t$,
- (T-2) for every edge $v_1 v_2 \in E_G$ there exists V_t such that $v_1, v_2 \in V_t$,
- (T-3) if there is a path $t_1 \dots t_2 \dots t_3$ in T then $V_{t_1} \cap V_{t_3} \subseteq V_{t_2}$.

Conditions (T-1) and (T-2) say the the graph G is the union of subgraphs induced by the set of vertices V_t ; the sets $V_t, t \in T$ as well as the subgraphs induced by $V_t, t \in T$ are called the *bags* of the tree decomposition. Condition (T-3) states that the bags of the tree decomposition are organized into a tree like fashion.

One of the most important feature of the tree decomposition concept is that it shows a natural correspondence between the properties of the separations of the graph and its tree decomposition:

Lemma 2.7. *Let G be a graph and let $(T, (V_t)_{t \in T})$ be a tree decomposition of G . Given an edge $t_1 t_2$ of T and let T_1, T_2 be the components of $T \setminus t_1 t_2$ such that $t_1 \in T_1$ and $t_2 \in T_2$.*

Then $V_{t_1} \cap V_{t_2}$ separates $U_1 := \bigcup_{t \in T_1} V_t$ from $U_2 := \bigcup_{t \in T_2} V_t$ in G .

Proof. Since T is a tree, then every t - t' path in T with $t \in T_1$ and $t' \in T_2$ contains both t_1 and t_2 . Therefore, it follows by (T-3) that $U_1 \cap U_2 \subseteq V_{t_1} \cap V_{t_2}$. To accomplish the proof it remains to show that G does not contain an edge

$u_1 u_2$ with $u_1 \in U_1 \setminus U_2$ and $u_2 \in U_2 \setminus U_1$. If such an edge exists then it follows from (T-2) that there exists $t \in V_T$ such that $u_1, u_2 \in V_t$. By assumption, we have chosen u_1 in $U_1 \setminus U_2$ and hence $t \in T_1$. Also, we have chosen u_2 in $U_2 \setminus U_1$ and hence $t \in T_2$. This is a contradiction because by the hypothesis we have $T_1 \cap T_2 = \emptyset$. $\square$

A *bag*, denoted β_t , $t \in V_T$, is the subgraph of G induced by vertices V_t . A *torso*⁴, denoted τ_t , is the bag β_t where we add an edge vv' to the multiset of edges of β_t for each $v, v' \in V_t \cap V_{t'}$ such that $tt' \in E_T$. Observe that after adding such edges the torso may become a multigraph even if the starting graph is simple. However, we can split the edges of a torso into two sets, the set of the original edges which belongs to the bag, and the set of edges which we have added, the latter are called the *virtual* edges.

3 Tutte's Theorem

The following Theorem, known as *Tutte decomposition Theorem* – which provides a tree decomposition of 2-connected graphs into cycles, bonds and 3-connected components – will be our main working tool.

Theorem 3.1. [Tut66] *Every 2-connected graph has a tree decomposition $(T, (V_t)_{t \in T})$ such that $|V_t \cap V_{t'}| = 2$ for every $tt' \in E_T$, and moreover every torso is either a k -bond ($k \geq 3$), or 3-connected or a cycle. Conversely, every graph with such a tree decomposition is 2-connected. Furthermore such a decomposition is unique.*

The key idea behind Tutte's Theorem consists in considering a particular set of 2-separations of G , these are the 2-separations which are *compatible* with all the other 2-separations. Two separations (U_1, U_2) and (W_1, W_2) are compatible if we can find $i, j \in \{1, 2\}$ such that $U_i \subseteq W_j$ and $U_{3-j} \supseteq W_{3-j}$. It follows from the proofs of Tutte's Theorem presented in the literature⁵ that (U_1, U_2) is a separation which is compatible with all the other separations if and only if $U_1 \cap U_2$ is a hinge. Finally Tutte's decomposition arises from considering a tree decomposition $(T, (V_t)_{t \in T})$ such that $V_t \cap V_{t'}$ is a hinge for all $tt' \in E_T$.

The original proof of this Theorem can be found in [Tut66]. An extension of this Theorem to locally finite⁶ graphs is referenced in [DSS95]. A generalization of this Theorem to arbitrary infinite graphs can be found in [Ric04]. In the sequel we shall refer to such a decomposition as *Tutte's decomposition*.

An example of a graph and its decomposition tree is depicted in Figure 3.1. Let $U_1 := V_G \setminus \{v_9\}$ and $U_2 := \{v_6, v_8, v_9\}$. Observe that (U_1, U_2) is a 2-separation of G . The set $U_1 \cap U_2 = \{v_6, v_8\}$ is a 2-separator of G but it is not a hinge because neither the subgraph $G[U_1]$ nor $G[U_2]$ is 2-connected. By inspecting the set of 2-separators, we can deduce that the set of hinges

⁴a torso and not a torsor. The plural of torso is torsos. We adopt the terminology of [Die05].

⁵For instance that of [Ric04], which uses the notion of hinges, and of [Die05], that uses the notion of compatible separations.

⁶An infinite graph is locally finite if the degree of each vertex is finite.

is $\{\{v_1, v_2\}, \{v_3, v_4\}, \{v_5, v_6\}, \{v_8, v_9\}\}$. The virtual edges arising from this decomposition are represented by dashed lines.

The following Lemma provides some useful Properties of Tutte's Tree.

Lemma 3.2. *Let G be a 2-connected graph and T its Tutte's tree. Then,*

(a) *T is a bipartite graph (T_1, T_2) where $t \in T_1$ if and only if V_t is a hinge.*

(b) *Let $t_1 \dots t_n$ be a path in T then, if h_i denotes the hinge V_{t_i} then*

(b.1) *for all $i = 1, \dots, n-1$, we have either $h_i \cap h_{i+1} = \emptyset$ or $|h_i \cap h_{i+1}| = 1$,*

(b.2) *if $h_p \cap h_q = \emptyset$ where $1 \leq p < q \leq n$, then for all $i \leq p$ we have that $h_i \cap h_q = \emptyset$, and*

(b.3) *if $h_p \cap h_q = \emptyset$ where $1 \leq p < q \leq n$, then for all $i \geq q$ we have that $h_p \cap h_i = \emptyset$.*

Proof. The statement (a) follows from Tutte's Theorem 3.1. The statements (b.1), (b.2), and (b.3) are a direct consequence of Lemma 2.7. $\square$

Lemma 3.3. [Ric04]. *Let G be 2-connected graph. A 2-separator $\{x, y\}$ is a hinge in G if and only if (i) either there are at least three $[x, y]$ -bridges, or (ii) there are two $[x, y]$ -bridges at least one of them is 2-connected.*

Tutte's decomposition tree of a given 2-connected graph gives rise to an algebraic expression in terms of k -bonds, cycles and 3-connected components and the 2-Sum operator. The latter operation consists in taking two graphs, choosing a 2-clique from each, identifying the vertices in the cliques and deleting the edges of the cliques, see Figure 3.2.

The k -sum operators on graphs, for $k = 1, 2, 3$,⁷ have been introduced in [Wag37] in the aim to prove a theorem which states that a graph which does not contain a K_5 as a minor may be expressed by means of these operations starting with the class of planar graphs and a particular graph on 8 vertices⁸.

Definition 3.4. Let G_1 and G_2 be graphs and let $e_1 = (a_1, b_1) \in E_{G_1}$ and $e_2 = (a_2, b_2) \in E_{G_2}$. Define the 2-Sum of G_1 and G_2 on e_1 and e_2 respectively, denoted $G_1 +_{e_1 e_2} G_2$, to be the graph obtained from the union of G_1 and G_2 by identifying the vertex a_1 with a_2 and the vertex b_1 with b_2 and deleting the edges e_1 and e_2 , see Figure 3.2.

It is clear that (see [Tut66]):

Lemma 3.5. *G and H are 2-connected if and only if $G +_{e_1 e_2} H$ is 2-connected, for each $e_1 \in E_G, e_2 \in E_H$.*

Note that the 2-bond is the neutral element of the 2-Sum. From the previous Lemma it follows that:

⁷The k -sum, for $k = 3$ is defined in a similar way of the 2-Sum operator apart that we take a 3-click of each graph.

⁸This Theorem has been extended to matroids in [Sey81] by extending these operations to matroids as well.




Figure 3.1: A graph with its Tutte's decomposition tree




Figure 3.2: The 2-Sum operator

Proposition 3.6. *Let $\mathcal{B}_2$ be the least class of graphs containing the k -bonds ($k \geq 3$), the cycles, the 3-connected graphs and closed under the 2-Sum operator. Then, $G \in \mathcal{B}_2$ if and only if G is 2-connected.*

Lemma 3.7. *Let G be 2-connected and let $(T, (V_t)_{t \in T})$ be its Tutte decomposition, then every torso τ of T is a minor of G .*

Proof. Let $t \in T$ and let $t_1, \dots, t_n$ be the neighbours of t in T . Let $T_i, i = 1, \dots, n$ be the component of $T \setminus tt_i$ that contains t_i , and let $\bar{T}_i$ be the complement of T_i w.r.t T . Let also $U_i := \cup_{t \in T_i} V_t$. Recall that if $U \subseteq V_G$ then the subgraph of G induced by U is denoted by $G[U]$.

We shall prove that the torso β_t is a minor of G . On the one hand, since the subgraph $G[U_i]$ is connected, see Theorem IV.20 of [Tut01], then there exists an u_i - w_i path π_i in $G[U_i]$ such that $\{u_i, w_i\} = V_t \cap V_{t_i}$. On the other hand, since $\{u_i, w_i\}$ is a 2-separator in G of U_i from $\bar{U}_i$ by Lemma 2.7, then $\{u_i, w_i\}$ is also a 2-separator in G of U_i from U_j where $i \neq j$ because $U_j \subseteq \bar{U}_i$. Therefore, the u_i - w_i paths, for $i = 1, \dots, n$, are independent. Observe that the graph β_t^+ consisting of the bag β_t where we add the u_i - w_i paths is a subgraph of G . Since the u_i - w_i paths are independent we can contract each path until we get a single edge and moreover the resulting graph is just the torsos τ_t . Hence τ_t is a minor of β_t^+ , and β_t^+ is a subgraph of G , hence τ_t is a minor of G . $\square$

From the previous Lemma we get the following particular case:

Lemma 3.8. *Let G be a 2-connected graph, such that $G = G_1 +_{e_1 e_2} G_2$. Then, $G_i + e_i$ is a minor of G , $i = 1, 2$.*

4 Entanglement, connectivity, and edge covering

The entanglement of a finite digraph G , denoted $\mathcal{E}(G)$, was defined in [BG05] by means of some games $\mathcal{E}(G, k)$, $k = 0, \dots, |V_G|$. The game $\mathcal{E}(G, k)$ is played on the digraph G by Thief against Cops, a team of k cops. The rules are as follows. Initially all the cops are placed outside the digraph, Thief selects and occupies an initial vertex of G . After Thief's move, Cops may do nothing, may place a

cop from outside the digraph onto the vertex currently occupied by Thief, may move a cop already on the digraph to the current vertex. In turn Thief must choose an edge outgoing from the current vertex whose target is not already occupied by some cop and move there. If no such edge exists, then Thief is caught and Cops win. Thief wins if he is never caught. The entanglement of G is the least $k \in \mathbb{N}$ such that k cops have a strategy to catch the thief on G . It will be useful to formalize these notions.

Definition 4.1. The entanglement game $\mathcal{E}(G, k)$ of a digraph G is defined by:

- Its positions are of the form (v, C, P) , where $v \in V_G$, $C \subseteq V_G$ and $|C| \leq k$, $P \in \{\text{Cops}, \text{Thief}\}$.
- Initially Thief chooses $v_0 \in V_G$ and moves to $(v_0, \emptyset, \text{Cops})$.
- Cops can move from (v, C, Cops) to (v, C', Thief) where C' can be
 - C : Cops skip,
 - $C \cup \{v\}$: Cops add a new Cop on the current position,
 - $(C \setminus \{x\}) \cup \{v\}$: Cops move a placed Cop to the current position.
- Thief can move from (v, C, Thief) to (v', C, Cops) if $(v, v') \in E_G$ and $v' \notin C$.

Every finite play is a win for Cops, and every infinite play is a win for Thief.

The following Proposition provides a useful variant of entanglement games, see [Bel08, §6].

Proposition 4.2. *Let $\tilde{\mathcal{E}}(G, k)$ be the game played as the game $\mathcal{E}(G, k)$ apart that Cops is allowed to retire a number of cops placed on the graph. That is, Cops moves are of the form*

- $(g, C, \text{Cops}) \rightarrow (g, C', \text{Thief})$ (generalized skip move),
- $(g, C, \text{Cops}) \rightarrow (g, C' \cup \{g\}, \text{Thief})$ (generalized replace move),

where in both cases $C' \subseteq C$. Then Cops has a winning strategy in $\mathcal{E}(G, k)$ if and only if he has a winning strategy in $\tilde{\mathcal{E}}(G, k)$.

We state one of the fundamental properties of the undirected graphs of bounded entanglement [Bel08, §6]:

Theorem 4.3. *The class of undirected graphs of entanglement at most n , for arbitrary fixed $n \in \mathbb{N}$, is minor closed, that is if G is minor of H then $\mathcal{E}(G) \leq \mathcal{E}(H)$.*

On the one hand, atoms of Tutte's decomposition are cycles and 3-connected components, on the other hand cycles have entanglement at most 3 [BS07]. The aim now is to investigate the structure of 3-connected components whenever the starting graph has entanglement 3. To this goal, we first establish the relation between the three notions of entanglement, connectivity and edge covering.

Lemma 4.4. *Let G be a k -connected graph with $|V_G| \geq k + 1$, then*

$$k \leq \mathcal{E}(G) \leq \mathcal{C}(G)$$

Proof. Let us prove first the inequality $\mathcal{E}(G) \leq \mathcal{C}(G)$. Let X be an edge cover set of G , we shall show that Cops have a winning strategy in $\mathcal{E}(G, |X|)$. While Thief is moving on a path π , Cops strategy consists in placing a cop on a vertex v of π if and only if v belongs to X , showing that the number of cops placed on the graph is at most $|X|$. If there is an infinite play then this implies that there is a cycle whose vertices are not covered by X , meaning that X is not an edge cover set of G , this is a contradiction. Therefore, Cops's strategy is winning and hence $\mathcal{E}(G) \leq \mathcal{C}(G)$.

To prove the inequality $k \leq \mathcal{E}(G)$, we use again Menger's Theorem 2.3, stating that *a graph is k -connected if and only if it contains k independent paths between any two vertices.*

We shall prove that Thief has a winning strategy in $\mathcal{E}(G, k - 1)$. To this goal it is sufficient to show the following conditions to hold: whenever $k - 1$ cops are placed on the graph, then

- (i) there is at least an edge whose both ends are not occupied by a cop. This is a consequence of the inequalities $\mathcal{C}(G) \geq k > k - 1$. The inequality $\mathcal{C}(G) \geq k$ is justified by the fact that every edge cover set of the graph G would disconnect it.
- (ii) when it is Thief's turn to move from some vertex v , then by Menger's Theorem, it follows that there is a free path (i.e. its vertices are not occupied by a cop) from v to some vertex w such that $ww' \in E_G$ and both w and w' are not occupied by a cop.

It follows that Thief's strategy consists in looking for an edge which is not occupied by a cop, choosing a free path to it, use this path to reach this edge, and iterating moves on it until Cops place a cop on one of its end points. This strategy can be iterated infinitely often, hence its a winning strategy for Thief in $\mathcal{E}(G, k - 1)$. $\square$

Lemma 4.5. *Let G be k -connected with $|V_G| \geq k + 1$. If $\mathcal{E}(G) = k$, then*

$$\mathcal{C}(G) = \mathcal{E}(G) = k.$$

Proof. We have already mentioned that $\mathcal{C}(G) \geq \mathcal{E}(G) = k$. To prove $\mathcal{C}(G) = k$ we assume that $\mathcal{C}(G) > k$ and we shall deduce a contradiction: we shall prove that Thief has a winning strategy in the game $\mathcal{E}(G, k)$.

We distinguish two cases in the game $\mathcal{E}(G, k)$ according to the number of cops placed on the graph:

Case (i). If the number of cops placed on the graph is at most $k - 1$ then the same conditions (i) and (ii) provided in the proof of Lemma 4.4 still hold, hence in this case Thief will never be caught.

Case (ii). If k cops are placed on the graph then consider the first position of the game for which the number of cops placed on the graph increases from

$k - 1$ to k , that is we consider the first Cops' add move $(v, C', Cops) \rightarrow (v, C' \cup \{v\}, Thief)$ where $|C' \cup \{v\}| = k$.

On the one hand, since $\mathcal{C}(G) > k$ by assumption then there exists an edge ww' which is not covered. On the other hand, since a cop is posted on v and $k - 1$ cops are posted on the remaining vertices, then by Menger's Theorem there are k independent paths linking v to w , therefore there exists a free path π from v to w . Thief's strategy consists in going from v to w through π and iterating moves on the edge ww' until Cops place a cop on either w or w' , say w , giving rise to a position of the form $(w, C, Thief)$ where $|C| = k$ and $w \in C$, returning back to the initial configuration. From the latter position Thief uses the same strategy described so far. Such a strategy can be iterated infinitely often, hence Thief has a winning strategy in $\mathcal{E}(G, k)$. This contradicts the hypothesis $\mathcal{E}(G) = k$. $\square$

5 The k -molecules

In this section, basing on Lemma 4.5, we shall characterize the structure of the k -connected graphs of entanglement k , for arbitrary $k \geq 1$.

Definition 5.1. Let $k, h \geq 1$, $B_k = \{b_1, \dots, b_k\}$ and $\mathcal{B} \subseteq B_k \times B_k$. A k -molecule $\vartheta_{B_k}^{\mathcal{B}, h}$ is the graph $G = (V, E)$ such that

1. $V = B_k \cup \{v_1, \dots, v_h\}$,
2. $E = \mathcal{B} \cup \{v_i b_j, 1 \leq i \leq h, 1 \leq j \leq k\}$.
3. $h \geq k - k'$, where k' is the connectivity of the subgraph of G induced by B_k .

The set B_k is called the base of the k -molecule.

The 1-molecules are the *stars*, the 2-molecules have been discussed in [BS07], and the 3-molecules $\vartheta_{B_3}^{\mathcal{B}, h}$ are pictured (up to graph isomorphism) in Figure 5.1; since no confusion will arise we have substituted $\mathcal{B}$ by its cardinality $|\mathcal{B}|$ ⁹.

The following Definition formalizes the fact that a graph is a k -molecule.

Definition 5.2. Let G be a graph and $B \subset V_G$ with $|B| = k$. We say that the pair (G, A) is a k -pre-molecule if there exists a k -molecule $\vartheta_{B_k}^{\mathcal{B}, h}$ and a graph isomorphism $\psi : G \rightarrow \vartheta_{B_k}^{\mathcal{B}, h}$ sending B to B_k .

We say that a G is an abstract k -molecule if there exists $B \subset V_G$ with $|B| = k$ such that the pair (G, B) is a k -pre-molecule.

Lemma 5.3. *Let G be an abstract k -molecule. If there exist $A_k, B_k \subset V_G$ such that each pair (G, A_k) and (G, B_k) is a k -pre-molecule. Then,*

⁹This is possible without confusion when $k = 3$, because a graph on 3 vertices is completely determined (up to isomorphism) by the number of its edges.




Figure 5.1: The structure of 3-molecules $\vartheta_{B_3}^{|\mathcal{B}|,h}$.

4.i the three subgraphs of G induced by $A_k \setminus B_k$, $B_k \setminus A_k$ and $V_G \setminus (A_k \cup B_k)$ are all discrete,

4.ii the subgraphs of G induced by A_k and B_k are isomorphic.

Proof.

4.i Assume that $A_k \setminus B_k$ is not discrete i.e. there exists $a_1, a_2 \in A_k$ with $a_1 a_2 \in E_{G_1}$. If we consider the k -pre-molecule (G, B_k) then we observe that $a_1 a_2 \in V_G \setminus B_k$, this is a contradiction since the graph induced by $V_G \setminus B_k$ must be discrete by the definition of the k -molecules. We get that the subgraph induced by $B_k \setminus A_k$ is discrete by just dualizing the above proof. To argue that $V_G \setminus (A_k \cup B_k)$ is discrete observe that $V_G \setminus (A_k \cup B_k) \subseteq V_G \setminus A_k$, and since $V_G \setminus A_k$ is discrete by the definition of the k -molecules, then it follows that $V_G \setminus (A_k \cup B_k)$ is also discrete.

4.ii Consider the mapping $\psi : G \rightarrow G$ such that ψ is a bijection from $A_k \setminus B_k$ to $B_k \setminus A_k$, and it is the identity on both $(A_k \cap B_k)$ and $V_G \setminus (A_k \cup B_k)$. Since the subgraphs induced by $A_k \setminus B_k$, $B_k \setminus A_k$, and $V_G \setminus (A_k \cup B_k)$ are discrete by 4.i, then for all $v_1, v_2 \in A_k$ and for all $w_1, w_2 \in B_k$, $\psi(v_i) = \psi(w_i)$, $i = 1, 2$, if and only if $\psi(v_1 v_2) = \psi(w_1 w_2)$. Hence the subgraphs of G induced by A_k and B_k are isomorphic.

□

In order to compute the connectivity of the k -molecules, the following Lemma provides a construction of them and a upper bound of their connectivity.

Lemma 5.4. *Let G be the graph constructed as follows: out of a graph B and a set of vertices $\{v_1, \dots, v_h\}$ add an edge between each $v_i, i = 1, \dots, h$ and each b in B . If B is k' -connected, then the connectivity of G is at least*

$$\min(|V_B|, k' + h).$$

Proof. Let $m = \min(|V_B|, k' + h)$, we shall prove that G is m -connected, by proving that every two vertices x, y are linked by at least m disjoint paths. We split the proof in three cases.

Case (i). If $x, y \in V_B$, then there are k' disjoint path in the subgraph induced by V_B from x to y because the latter is k' -connected. Moreover there are h disjoint paths of the form $xv_1y, x_2y, \dots, xv_hy$ where $v_i \in V_G \setminus V_B$.

Case (ii). If $x, y \in V_G \setminus V_B$. In this case there are $|V_B|$ disjoint paths of the form $xb_1y, xb_2y, \dots, xb_ky$, where $b_i \in V_B$.

Case (iii). If $x \in V_B$ and $y \in V_G \setminus V_B$, then the $k' + h$ disjoint paths are $\Pi_1 \cup \Pi_2 \cup \Pi_3$ where :

- $\Pi_1 = \{xy\}$, recall that $xy \in E_G$ by definition.
- To exhibit Π_2 recall first that since the graph B is k' connected then x has at least k' neighbors in B , let $b_1, \dots, b_{k'}$ be such neighbors. Therefore we let $\Pi_2 = \{xb_1y, \dots, xb_{k'}y\}$.
- Finally,

$$\Pi_3 = \{xv_1b'_1y, \dots, xv_ib'_iy, \dots, xv_{h-1}b'_{h-1}y\}$$

where $\{b_1, \dots, b_{k'}\} \cap \{b'_1, \dots, b'_{h-1}\} = \emptyset$.

It is straightforward to check that paths in $\Pi_1 \cup \Pi_2 \cup \Pi_3$ are disjoint, they share just their two end points. Moreover $|\Pi_1 \cup \Pi_2 \cup \Pi_3| = 1 + k' + (h - 1) = k' + h$.

□

Now we state the main combinatorial properties of the k -molecules.

Proposition 5.5. *Let $G = \vartheta_{B_k}^{\mathcal{B}_k, h}$ be a k -molecule. Then*

1. *the connectivity of G is k ,*
2. *G has B_k as a minimal edge cover and hence $\mathcal{C}(G) = k$, and*
3. *the entanglement of G equals k .*

Proof.

1. Observe that if G is a clique then it should be a $(k + 1)$ -clique. Moreover, this holds if and only if the subgraph of G induced by $\mathcal{B}_k$ is a k -clique and $h = 1$. In this case the connectivity of G is k by Convention 2.5. Assume that G is not a clique. On the one hand, by lemma 5.4, it follows that the connectivity of G is at least $\min(|B_k|, k' + h) = \min(k, k' + h)$ and since $h \geq k - k'$ by the definition of the k -molecules, then $\min(k, k' + h) = k$, showing that the connectivity of G is at least k . On the other hand, we need the Claim:

Claim 5.6. *Let $G := \vartheta_{B_k}^{B,h}$ be a k -molecule. If G is not a clique then $h \geq 2$.*

Proof. We distinguish two cases according to the nature of $G[B_k]$.
 If $G[B_k]$ is a k -clique, then we need $h \geq 2$, because if $h = 1$ then G would be a $(k+1)$ -clique contradicting the hypothesis.
 If $G[B_k]$ is not a clique, then there exist at least two vertices $b, b' \in B_k$ such that $bb' \notin E_G$, therefore $B_k \setminus \{b, b'\}$ is a $(k-2)$ -separator of b from b' in $G[B_k]$. This implies that the connectivity of $G[B_k]$ is at most $k-2$. From the definition of the k -molecules, we have $h \geq k - k'$ where k' is the connectivity of $G[B_k]$, hence $h \geq k - (k-2) = 2$. This ends the proof of the Claim. $\square$

Since G is not a clique then, according to the Claim, we have $h \geq 2$ and hence B_k is a k -separator in G of any two vertices in $V_G \setminus B_k$ and in this case G can not be $(k+1)$ -connected. We conclude that the connectivity of G is k .

2. Since G is k -connected and $|V_G| \geq k+1$ then by Lemma 4.4 we get $\mathcal{C}(G) \geq k$. It is easy to check that B_k is an edge cover of G , hence $\mathcal{C}(G) \leq |B_k| = k$. Therefore $\mathcal{C}(G) = k$.
3. On the one hand, since G is k -connected then from Lemma 4.4 we obtain $\mathcal{E}(G) \geq k$. On the other hand, from the same Lemma 4.4, we have $\mathcal{E}(G) \leq \mathcal{C}(G)$ and from the previous item we got $\mathcal{C}(G) = k$, thus $\mathcal{E}(G) \leq k$. We conclude that $\mathcal{E}(G) = k$.

$\square$

So far we have stated and proved the main properties of the k -molecules. Conversely, the following Proposition characterizes the graphs for which the entanglement, the connectivity and the cyclicity coincide.

Proposition 5.7. *If G is k -connected with $\mathcal{E}(G) = k$, then G is an abstract k -molecule.*

Proof. Let G be a graph as stated. Since $\mathcal{E}(G) = k$ and G is k -connected then by Lemma 4.5 it follows that $\mathcal{E}(G) = \mathcal{C}(G) = k$, thus let $B_k = \{b_1, \dots, b_k\}$ be a minimal edge cover set of G and let k' be the connectivity of $G[B_k]$.
 We claim that vertices in $V_G \setminus B_k$ are at distance one from B_k because B_k is an edge cover set of G . For the same reason the subgraph of G induced by $V_G \setminus B_k$ is discrete. Thus, for each $v \in V_G \setminus B_k$ there exists $b \in B_k$ such that $vb \in E_G$. Let $v \in V_G \setminus B_k$ and $\mathcal{N}_v$ be the set of its neighbors, clearly $\mathcal{N}_v \subseteq B_k$ because $G[V_G \setminus B_k]$ is discrete. If $|\mathcal{N}_v| < k$ meaning that $\mathcal{N}_v \subsetneq B_k$ then clearly $\mathcal{N}_v$ separates v from $B_k \setminus \mathcal{N}_v$. This contradicts the assumption that G is k -connected. We conclude that $\mathcal{N}_v = B_k$ for each $v \in V_G \setminus B_k$.
 Finally, to accomplish the proof that G , coming with the desired data, is a k -molecule, it remains just to show that $|V_G \setminus B_k| \geq k - k'$ where k' is the




Figure 5.2: Two possible ways to view a 2-molecule.

connectivity of $G[B_k]$. Towards a contradiction, assume that $|V_G \setminus B_k| < k - k'$. Since the connectivity of $G[B_k]$ is k' then there exists a k' -separator in $G[B_k]$, let $S_{k'}$ be such a separator and assume that it separates b_1 from b_2 ; $b_1, b_2 \in B_k$. Therefore $S_{k'} \cup (V_G \setminus B_k)$ separates also b_1 from b_2 . A simple computation shows that $|S_{k'} \cup (V_G \setminus B_k)| < k$, contradicting the fact that G is k -connected. We conclude that the pair (G, B_k) is a k -premolecule, moreover G may be written as $\vartheta_{B_k}^{\mathcal{B}, h}$ where $\mathcal{B}$ are the edges of the subgraph of G induced by B_k , and $h = |V_G \setminus B_k|$. $\square$

5.1 Classification of the 3-molecules

A natural question arises: in how many manners a k -molecule may be written? To illustrate this question consider the graph G depicted in Figure 5.2. If we take $A_2 = \{a, b\}$, then the graph G may be viewed as the two molecule $\vartheta_{A_2}^{\emptyset, 2}$. If we take $B_2 = \{c, d\}$, then G may be considered as the 2-molecule $\vartheta_{B_2}^{\emptyset, 2}$.

A k -molecule may admit many bases giving rise to what we call *ambiguous* molecules. The formal definition of ambiguity follows.

Definition 5.8. Let G be an abstract k -molecule. We say that G is non ambiguous if there exists just one set $B \subset V_G$ such that the pair (G, B) is a k -pre-molecule. Similarly, a k -molecule ϑ is non ambiguous if ϑ viewed as a graph is non ambiguous.

The following Proposition gives an explicit characterization the class of ambiguous 3-molecules.

Proposition 5.9. Let ϑ be a 3 molecules $\vartheta_{B_3}^{\mathcal{B}, h}$, then ϑ is ambiguous if and only if one of the following cases holds:

- I. $|\mathcal{B}| = 3$ and $h = 1$, that is ϑ is the 4-clique,
- II. $|\mathcal{B}| = 2$ and $h = 2$,

III. $|\mathcal{B}| = 0$ and $h = 3$, that is ϑ is the complete bipartite graph $K_{3,3}$.

These graphs are depicted in figure 5.3




Figure 5.3: The set of ambiguous 3-molecules.

Proof. First, we give a useful property of ambiguous k -molecules:

Claim 5.10. *If G is an ambiguous k -molecule, and A_k, B_k are two different bases of G , then $A_k \cup B_k = V_G$.*

Proof. Assume that $A_k \cup B_k \subsetneq V_G$ and then let $w \in V_G \setminus (A_k \cup B_k)$. Hence, for each $b \in B_k$ we have by the definition of the k -molecules that $wb \in E_G$. This implies that the subgraph of G induced by $V_G \setminus A_k$ is not discrete, contradicting the definition of the k -molecules. We conclude that $A_k \cup B_k = V_G$. This ends the proof of Claim. $\square$

Let $G := \vartheta_{B_3}^{\mathcal{B},h}$ be an ambiguous 3-molecule and A_3, B_3 be two distinct bases of G , thus by the previous Claim we get $A_3 \cup B_3 = V_G$. We recall first that Lemma 5.3 states that both the graphs induced by $A_3 \setminus B_3$ and $B_3 \setminus A_3$ are discrete and moreover the graphs induced by A_3 and B_3 are isomorphic. We distinguish three cases according to $|A_3 \cap B_3|$.

1. $|A_3 \cap B_3| = 0$. Since both $G[A_3]$ and $G[B_3]$ are discrete, then G is the complete bipartite graph $K_{3,3}$, i.e. G is the 3-molecule $\vartheta_{B_3}^{\mathcal{B},3}$ where $\mathcal{B} = 0$.
2. $|A_3 \cap B_3| = 1$. Let $A_3 = \{w, a_1, a_2\}$ and $B_3 = \{w, b_1, b_2\}$. First, $a_1a_2 \notin E_G$ because $G[A_3 \setminus B_3]$ is discrete, and also $b_1b_2 \notin E_G$ because $G[B_3 \setminus A_3]$ is discrete. Second, we shall argue that $\{wa_i, wb_i, i = 1, 2\} \subset E_G$. Assume that $wa_1 \notin E_G$, then $\{b_1, b_2\}$ is a 2-separator in G of a_1 from a_2 . This is a contradiction because the 3-molecules are 3-connected. We deduce that $wa_1 \in E_G$. By symmetry, we obtain also that $wa_2, wb_1, wb_2 \in E_G$. We conclude that, in this case, G is the 3-molecule $\vartheta_{B_3}^{\mathcal{B},2}$ where $\mathcal{B} = 2$.

3. $|A_3 \cap B_3| = 2$. Observe that in this case $h = 1$ in $\vartheta_{B_3}^{\mathcal{B},h}$. We shall argue next that both $G[A_3]$ and $G[B_3]$ are the 3-clique. Recall that in the k -molecule $\vartheta_{B_k}^{\mathcal{B},h}$ we have $h \geq k - k'$ where k' is the connectivity of $G[B_k]$. If $|E_{G[A_3]}| \leq 2$, then the connectivity of $G[A_3]$ is at most 1 and hence in the 3-molecule $\vartheta_{A_3}^{\mathcal{B},h}$ we should have $h \geq k - k' \geq 3 - 1 = 2$, this is a contradiction because we have already mentioned that $h = 1$. We conclude that $|E_{G[A_3]}| = 3$ meaning that G is the 4-clique, i.e. the 3-molecule $\vartheta_{B_3}^{\mathcal{B},1}$ where $|\mathcal{B}| = 3$.

□

6 Tree decomposition of graphs of entanglement at most 3

The main result of [BS07] states that a graph of entanglement 2 has a tree decomposition into 2-connected components such that the latter are the 2-molecules that come with a prescribed set of articulation points. Conversely, starting with the 2-molecules and the 1-Sum operator we have been able to generate the class of graphs of entanglement 2. Now we shall follow this approach to deal with the class of graphs of entanglement 3.

We find some *necessary* conditions on the structure of the Tutte's tree to be a tree decomposition of a 2-connected graph of entanglement 3. The necessary conditions deal with three features of the tree: (i) conditions on the structure of the 3-connected components: they are the 3-molecules. This is direct consequence of the results proved in the previous sections, (ii) conditions on the hinges are given in a similar way of those given on the articulation points when the starting graph has entanglement 2, and (iii) conditions on the diameter of the tree.

6.1 Necessary conditions on Tutte's tree

Since a 2-connected graph may be written by means of the 2-Sum operator, we begin by inspecting the main cases for which the 2-Sum operator increases the entanglement¹⁰. In the next, the symbol $+$ denotes the 2-Sum operator on 2-connected graphs given in Definition 3.4.

From now on we shall deal particularly with the 3-molecules and write $\vartheta_{a,b,c}^{\mathcal{B},h}$ instead of $\vartheta_{\{a,b,c\}}^{\mathcal{B},h}$.

Lemma 6.1. *Let $\vartheta_{a,b,c}^{\mathcal{B},h}$ be a 3-molecule and C_3 be the 3-cycle on the vertices $\{a, v_1, z\}$. Define the graph G as follows:*

$$G = C_3 +_{av_1} \vartheta_{a,b,c}^{\mathcal{B},h}.$$

¹⁰This idea has been already considered in [BS07] where we have looked for the cases for which the 1-sum operator increases the entanglement. There, the 1-sum operator which does not increase the entanglement is called the legal 1-Sum.

If $\{a, v_1\}$ does not belong to any minimal edge cover set of $\vartheta_{a,b,c}^{\mathcal{B},h}$, then $\mathcal{E}(G) \geq 4$

Proof. The graph G may be viewed as the graph that results from inserting the new vertex z in the edge av_1 of $\vartheta_{a,b,c}^{\mathcal{B},h}$. Let us abbreviate $\vartheta_{a,b,c}^{\mathcal{B},h}$ by ϑ . Define $f : G \rightarrow \vartheta$ as follows: $f(v) = v$ if $v \neq z$, and $f(z) = a$. Thief's strategy in $\tilde{\mathcal{E}}(\vartheta, 3)$ ¹¹ (even if it is losing) that forces Cops to occupy a minimal edge cover set of ϑ , this strategy exists by Lemma ??, will allow us to construct a winning strategy for Thief in $\mathcal{E}(G, 3)$. Every position (v, C_ϑ, P) of $\tilde{\mathcal{E}}(\vartheta, 3)$ is matched with a position (g, C_G, P) of $\mathcal{E}(G, 3)$ such that the following conditions hold:

- $f(g) = w$ and $f(C_G) = C_\vartheta$. (COPS)
- if Thief moves from $(a, C_\vartheta, Thief)$ to $(v_1, C_\vartheta, Cops)$,
then $a \in C_\vartheta$. (COP-ON- a)

The condition (COPS) states essentially that the cops in $\tilde{\mathcal{E}}(\vartheta, 3)$ are placed on the image of cops in $\mathcal{E}(G, 3)$ by the function f defined above. The condition (COP-ON- a) states that whenever Thief leaves vertex a to v_1 then a cop must already be placed on a .

A Thief's move $M_\vartheta = (u, C_\vartheta, Thief) \rightarrow (w, C_\vartheta, Cops)$ in $\tilde{\mathcal{E}}(\vartheta, 3)$ is simulated either by a move or a sequence of moves in $\mathcal{E}(G, k)$ according to the edge uw .

1. If $uw \neq v_1a$, then the move M_G is simulated in $\mathcal{E}(G, 3)$ by the same Thief's move. Observe that the latter move is possible in $\mathcal{E}(G, 3)$ because $w \in C_G$ then already $f(w) = w \in f(C_G) = C_\vartheta$, and this is impossible.
2. If $(u, w) = (v_1, a)$, then the move M_G is simulated in $\mathcal{E}(G, 3)$ by the following sequence of moves:

$$(v_1, C_G, Thief) \rightarrow (z, C_G, Cops) \rightarrow (z, C'_G, Thief) \rightarrow (a, C'_G, Cops)$$

This sequence is possible. If Thief can not perform such moves then either $z \in C_G$ or $a \in C'_G$. If $z \in C_G$ then $f(z) = a \in f(C_G) = C_\vartheta$, which is impossible. If $a \in C'_G$, then already $a \in C_G$ and hence $f(a) = a \in f(C_G) = C_\vartheta$, which is also impossible. Observe that in this case, Thief's move in $\tilde{\mathcal{E}}(\vartheta, 3)$ is simulated by a sequence of Thief's moves in $\mathcal{E}(G, 3)$, and the latter are interleaved with Cops moves, and then the position of Cops in $\tilde{\mathcal{E}}(\vartheta, 3)$ such be updated using the function f . So, it remains to show that Cops related moves in $\tilde{\mathcal{E}}(\vartheta, 3)$ respect the rule of the game.

From the latter position in $\mathcal{E}(G, 3)$, Cops next move is a from $(a, C'_G, Cops) \rightarrow (a, C''_G, Thief)$. Hence in $\tilde{\mathcal{E}}(\vartheta, 3)$ Cops related move should be of the form $(a, C_\vartheta, Cops) \rightarrow (a, f(C''_G), Thief)$. Let us compute $f(C''_G)$ in term of C_ϑ . Observe first that $C''_G = (C_G \setminus A) \cup B$ where $\emptyset \subseteq B \subseteq \{z, a\}$ and $A \subset C_G$

¹¹The game $\tilde{\mathcal{E}}(G, k)$ is defined as the game $\mathcal{E}(G, k)$ apart that Cops can retire a number of cops, the two versions are equivalent, Proposition 4.2.

with $|A| \leq 2$.

$$\begin{aligned} f(C_G'') &= f[(C_G \setminus A) \cup B] \\ &= f(C_G \setminus A) \cup f(B) \\ &= [f(C_G) \setminus f(A)] \cup f(B) \end{aligned}$$

where $\emptyset \subseteq Z \subseteq \{z\}$ and $f(B) \subseteq \{z, a\}$. Therefore,

$$\begin{aligned} f(C_G'') &= (f(C_G) \setminus f(A)) \cup (Z \cup f(B)) \\ &= (C_\vartheta \setminus f(A)) \cup Z' \end{aligned}$$

On the one hand $A \subset C_G$ and hence $f(A) \subset f(C_G) = C_\vartheta$. On the other hand $Z' = Z \cup f(B) \subseteq \{z\} \cup \{z, a\} = \{z, a\}$. We conclude that Cops' move under discussion respects the rules of the game.

3. If $(u, w) = (a, v_1)$, then the move M_ϑ is simulated in $\mathcal{E}(G, 3)$ by Thuef's iteration on az until a cop is placed either on a or z and then Thief goes to v_1 . That is, it is the following sequence:

$$\begin{aligned} M_G^* &= (a, C_G, Thief) \rightarrow (z, C_G, Cops) \rightarrow (z, C_G, Thief) \rightarrow (a, C_G, Cops) \\ &\rightarrow (a, C_G, Thief) \rightarrow (z, C_G, Cops) \\ &\rightarrow \dots \\ &\rightarrow (a, C_G, Cops) \rightarrow (a, C_G', Thief) \\ &\rightarrow (z, C_G', Cops) \rightarrow (z, C_G'', Thief) \\ &\rightarrow (v_1, C_G'', Cops) \end{aligned}$$

Such that $C_G' \neq C_G$ or $C_G'' \neq C_G$. Let us show that Thief can perform such moves, i.e. $z \notin C_G$ and $v_1 \notin C_G''$.

If $v_1 \in C_G''$ then $v_1 \in C_G$, and hence $f(v_1) = v_1 \in f(C_G) = C_\vartheta$, this is impossible.

If $z \in C_G$ the $f(z) = a \in f(C_G) = C_\vartheta$, then let us come back to the previous round of simulation. We mean if we consider Thief's previous moves in $\tilde{\mathcal{E}}(\vartheta, 3)$, then they are of the form

$$(a^{-1}, C_\vartheta^{-1}, Thief) \rightarrow (a, C_\vartheta^{-1}, Cops) \rightarrow (a, C_\vartheta, Thief)$$

and since we have supposed that $z \in C_G$, then $f(z) = a \in f(C_G) = C_\vartheta$ and moreover $a \in C_\vartheta^{-1}$, which is impossible.

Let us argue now that Cops' next move in $\tilde{\mathcal{E}}(\vartheta, 3)$ respects the rules of the game. From the latter position in $\mathcal{E}(G, 3)$, Cops' next move would be of the form $(v_1, C_G'', Cops) \rightarrow (v_1, C_G^*, Thief)$, and hence Cops' in $\tilde{\mathcal{E}}(\vartheta, 3)$ would reply, according to condition (COPS) by the move $(v_1, C_\vartheta, Cops) \rightarrow (v_1, f(C_G^*), Thief)$. Observe first that $a \in C_\vartheta$ by the condition (COP-ON- a). In order to show that the latter move respect the rules of the game, we compute $f(C_G^*)$ in term of C_ϑ . Note that $C_G^* = (C_G \setminus A) \cup B$ where

$\emptyset \subseteq B \subseteq \{a, z, v_1\}$ and $A \subseteq C_G$. As in the previous case, a simple computation shows that $f(C_G^*) = (C_\vartheta \setminus f(A)) \cup f(B')$ where $B' \subseteq \{a, z, v_1\}$. On the one hand $A \subseteq C_G$ implying $f(A) \subseteq f(C_G) = C_\vartheta$. On the other hand $f(B') \subseteq \{a, v_1\}$. However, we have mentioned that $a \in C_\vartheta$, and hence $f(C_G^*) = (C_\vartheta \setminus f(A)) \cup B''$ where $B'' \subseteq \{v_1\}$. This shows that Cops' move in question respects the rules of the game.

So far we have described the simulation between the games $\mathcal{E}(G, 3)$ and $\tilde{\mathcal{E}}(\vartheta, 3)$ and shown that it is consistent. Now we shall show that the hypothesis of the Lemma under proof imply implicitly some restrictions on the 3-molecule $\vartheta_{a,b,c}^{\mathcal{B},h}$ provided in this Lemma.

Claim 6.2. *the 3-molecule $\vartheta_{a,b,c}^{\mathcal{B},h}$ described in Lemma 6.1 is not the 4-clique, and hence $h \geq 2$.*

Proof. If $\vartheta := \vartheta_{a,b,c}^{\mathcal{B},h}$ is the 4-clique, then any set of vertices of size 3 forms a minimal edge cover of ϑ , implying that a, v_1 belongs to some minimal edge cover of ϑ , contradicting the hypothesis. This ends the proof of the Claim. $\square$

Assume that ϑ is ambiguous and let $\{a', b', c'\}$ be an other minimal edge cover of ϑ . Since the edge $v_1 a$, where the 2-Sum is performed, does not belong to any minimal edge cover of ϑ , then it is invariant w.r.t. changing the bases of ϑ . This shows that we can deal with ϑ like wise $\{a, b, c\}$ is the unique base.

If Thief is trapped in $\tilde{\mathcal{E}}(\vartheta, 3)$ then we have a position of the form $(v, \{a, b, c\}, \text{Thief})$ where $v \in \{a, b, c\}$. The latter position is matched with a position (v, C_G, Thief) of $\mathcal{E}(G, 3)$ where $f(C_G) = \{a, b, c\}$. Therefore, either $C_G = \{a, b, c\}$ or $C_G = \{z, b, c\}$.

Case (i). If $C_G = \{a, b, c\}$, then Thief can go to v_1 and iterates moves on $v_1 z$ forcing Cops to put a cop on v_1 or z . If Cops put a cop on v_1 then the image of Cops on ϑ by f is no longer a minimal edge cover, and hence Thief plays in $\tilde{\mathcal{E}}(\vartheta, 3)$ with the strategy that consists in forcing Cops to occupy a minimal edge cover set of ϑ , i.e. $\{a, b, c\}$. If Cops put a cop on z , then this cop comes from a, b , or c .

If this cop comes from b or c , then the image of cops on ϑ by f is either $\{a, c\}$ or $\{a, b\}$ which is not a minimal edge cover set of ϑ , and hence Thief forces Cops to occupy again a minimal edge cover of ϑ .

If this cop comes from a then we go to *Case(ii)*.

Case (ii). If $C_G = \{z, b, c\}$, then assume that the current vertex occupied by Thief is denoted by $x \in \{z, b, c\}$, and hence Thief's can choose the path xav_2 and¹² iterates moves on $v_2 a$ forcing Cops to put a cop on a , coming back to *Case(i)*.

Such a strategy for Thief in $\mathcal{E}(G, 3)$ can be iterated infinitely often, that is, it is a winning strategy, and hence $\mathcal{E}(G) \geq 4$. This ends the proof of Lemma 6.1. $\square$

¹²The vertex v_2 exists because $h \geq 2$ in $\vartheta = \vartheta_{a,b,c}^{\mathcal{B},h}$

Corollary 6.3. *Let G be 2-connected such that $\mathcal{E}(G) = 3$ and T be its Tutte decomposition. Let $t_1 t_2 \in E_T$ such that the torso τ_{t_1} is 3-connected, then there exists $B_3 \subset V_{\tau_{t_1}}$ such that (i) the pair (τ_{t_1}, B_3) is a 3-premolecule, and (ii) $V_{t_1} \cap V_{t_2} \subset B_3$.*

Proof.

(i). Since τ_{t_1} is 3-connected, then by Lemma 4.4, we get $\mathcal{E}(\tau_{t_1}) \geq 3$. By Lemma 3.7, the torsos τ_{t_1} is a minor of G , hence $\mathcal{E}(\tau_{t_1}) \leq 3$. Hence, $\mathcal{E}(\tau_{t_1}) = 3$. Since τ_{t_1} is 3-connected and $\mathcal{E}(\tau_{t_1}) = 3$, then it follows from Lemma 5.7, that there exists $B_3 \subset V_{\tau_{t_1}}$ such the pair (τ_{t_1}, B_3) is a 3-premolecule.

(ii). We need the Claim:

Claim 6.4. *Let $\{v_1 v_2\} = V_{t_1} \cap V_{t_2}$, then the graph $\tau_{t_1} +_{v_1 v_2} C_3$ is a minor of G .*

The Claim implies that $\mathcal{E}(\tau_{t_1} +_{v_1 v_2} C_3) \leq 3$, then from Lemma 6.1 we get that v_1, v_2 should belong to a minimal edge cover of τ_{t_1} . Let B'_3 be such a minimal edge cover, then it follows that the pair (τ_{t_1}, B'_3) is again a 3-premolecule. $\square$

6.2 On the diameter of the tree decomposition.

Now we shall give conditions on the diameter of the tree decomposition if the graph has entanglement 3. To give an upper bound for the diameter of the tree, we have noticed that a domino of a prescribed length is a typical excluded minor. The *domino* D_n is the graph of vertices $V_{D_n} = \{v_i, w_i \mid i = 0, \dots, n\}$ and edges $E_{D_n} = \{v_i v_{i+1}, w_i w_{i+1} \mid i = 0, \dots, n-1\} \cup \{v_i w_i \mid i = 0, \dots, n\}$.

In the following, if n is even then we prefer that the set of vertices of the n -domino would be of the form

$$\{v_i, w_i \mid i = -\frac{n}{2}, \dots, 0, \dots, \frac{n}{2}\}$$

The 14-domino is depicted in Figure 6.2.




Figure 6.1: The 14-Domino D_{14}

Lemma 6.5. *The entanglement of the domino D_{14} is at least 4.*

Proof. We shall describe a winning strategy for Thief in $\mathcal{E}(D_{14}, 3)$. First, let us fix some terminology and notations. Let $V = \{v_i \mid i = -7, \dots, 7\}$ and $W = \{w_i \mid i = -7, \dots, 7\}$. Let $\mathcal{L}_4$, the left 4-domino, be the subgraph of D_{14}

induced by the vertices $\{v_i, w_i \mid i = -7, \dots, -3\}$. Similarly, we let $\mathcal{R}_4$, the right 4-domino, be the subgraph of D_{14} induced by the vertices $\{v_i, w_i \mid i = 3, \dots, 7\}$. We let also $\mathcal{C}_2$, the central 2-domino, be the subgraph induced by the vertices $\{v_i, w_i \mid i = -1, 0, 1\}$. If S is a sub-domino of D_{14} then the $S \cap V$ -path (resp. $S \cap W$ -path) is the path induced by vertices $V_S \cap V$ (resp. $V_S \cap W$). Before giving Thief's winning strategy, we describe it informally:

- Step 1. Thief plays on $\mathcal{C}_2$ forcing Cops to place 3 cops on it. According to the last position of Thief in $\mathcal{C}_2$, Thief chooses the left or the right 4-domino, and moreover chooses an extremal vertex of it among $\{v_i, w_i \mid i = -3, 3\}$. Let us assume that Thief chooses $\mathcal{L}_4$ and the vertex v_{-3} ,
- Step 2. from the choice $(v_{-3}, \mathcal{L}_4)$, Thief iterates moves on the 4-path $\mathcal{L}_4 \cap V$ starting from v_{-3} until 2 cops are placed on this path, and at this moment
 - 2.1. if there is no cop on $\mathcal{C}_2$, then Thief goes to $\mathcal{C}_2$, and then iterates the strategy from Step 1,
 - 2.2. if there is a cop on the $\mathcal{C}_2$, then there is no cop on $\mathcal{R}_4$, and hence Thief chooses a path to $\mathcal{R}_4$ and an extremal vertex of it among $\{v_3, w_3\}$, and then iterates the strategy, up to symmetry, from Step 2.

The formal proof is split into two parts. The first part, called the *intra-steps*, consists in proving that the strategy for Thief described beside each step is realizable. The second part, called the *inter-steps*, consists in proving that the passage between steps is possible. Technically speaking, the inter-steps proofs, are devoted to prove that the path - that leads to the desired sub-domino- is free of Cops.

Let us begin by Step 1. To argue that Thief can play on $\mathcal{C}_2$ in such a way he forces 3 cops to be placed on it, it is sufficient to observe that $\mathcal{E}(\mathcal{C}_2) > 2$ because $\mathcal{C}_2$ does not belong to the class ζ_2 of the graphs of entanglement 2, see [BS07]. The following Claim ensures that the passage from Step 1 to Step 2 is possible.

Claim 6.6. *Thief has a strategy to play in $\mathcal{C}_2$ in such a way, once 3 cops are placed on $\mathcal{C}_2$, he can go in a horizontal way either to $\mathcal{L}_4$ or $\mathcal{R}_4$.*

Proof. Assume that Thief is trapped in $\mathcal{C}_2$ without being able to find a path neither to $\mathcal{L}_4$ nor to $\mathcal{R}_4$. This implies that Thief is on v_0 or on w_0 - otherwise he is able to go to $\mathcal{L}_4$ or to $\mathcal{R}_4$ - and moreover he is surrounded by 3 Cops in such a way he can not move down, left and right. That gives rise to the final position $(v_0, \{v_{-1}, w_0, v_1\}, \text{Thief})$ ¹³. Coming back two moves before, we get a position of the form $(x, \{v_{-1}, w_0, v_1\}, \text{Thief})$ where $x \in \{v_{-1}, w_0, v_1\}$. From the latter position Thief is clearly able to go to either $\mathcal{L}_4$ or $\mathcal{R}_4$. This ends the proof of the Claim. $\square$

¹³Since we reason up to symmetry, we put the second position $(w_0, \{w_{-1}, v_0, w_1\}, \text{Thief})$ in the back ground.

Assume now that 3 cops are placed on $\mathcal{C}_2$ and Thief is on vertex v_{-1} . The other cases are handled by symmetry. It follows from the Claim that Thief is able to choose the pair $(v_{-3}, \mathcal{L}_4)$ by going through the path $v_{-1}v_{-2}v_{-3}$.

Let us prove that the strategy described in Step 2 is possible. Once the pair $(v_{-1}, \mathcal{L}_4)$ is chosen and Thief is on v_{-1} , then Thief restricts his moves on the $\mathcal{L}_4 \cap V$ -path, the latter is of length 4 and therefore it has entanglement 2. Hence Thief has a strategy to force 2 cops to be placed on this path. At this moment either there is a cop on $\mathcal{C}_2$ or not.

- If there is a cop on $\mathcal{C}_2$, and since 2 cops are placed on the $\mathcal{L}_4 \cap V$ path, then there is no cop on $\mathcal{R}_4$, and moreover there is a free path leading to one of its left extremal points from the current vertex i.e.
 - (a) if there is a cop on $\mathcal{C}_2 \cap W$, then Thief goes from $v_p \in \mathcal{L}_4 \cap V$ to v_3 through the free path $v_p w_p w_{p+1} \dots w_{-2} v_{-2} v_{-1} \dots v_3$. Therefore iterates the strategy from Step 2 out of the pair $(v_3, \mathcal{R}_4)$,
 - (b) if there is a cop on $\mathcal{C}_2 \cap V$, then Thief goes from $v_p \in \mathcal{L}_4 \cap V$ to w_3 through the free path $v_p w_p w_{p+1} \dots w_3$. Therefore, iterates the strategy from Step 2 out of the pair $(w_3, \mathcal{R}_4)$

Remark 6.7. We emphasize that in case (a), the path chosen by Thief does not pass through $(\mathcal{R}_4 \cap W) \cup \{w_2\}$, the latter vertices are indeed free of cops and hence they might be used later by Thief. Also, in the case (b), the path chosen by Thief does not pass through $(\mathcal{R}_4 \cap V) \cup \{v_2\}$, the latter vertices are free of cops, and they would be used later by Thief.

- if there is no cop on $\mathcal{C}_2$ then, from the current vertex $v_p \in \mathcal{L}_4 \cap V$ Thief goes to $\mathcal{C}_2$ through the path $v_p w_p w_{p+1} \dots w_{-1}$. The freeness of this path is ensured by Remark 6.7. From the latter position, iterates the strategy from Step 1

The strategy for Thief in $\mathcal{E}(D_{14}, 3)$ described so far can be iterated infinitely often, that is, it is a winning strategy for Thief. Therefore $\mathcal{E}(D_{14}) \geq 4$. $\square$

Given a graph G and its Tutte's decomposition tree T , we call the *spread* of a vertex $v \in V_G$, denoted by δ_v , the number of hinges which v belongs to; i.e.

$$\delta_v = |\{ \{v, x\} \text{ s.t. } \{v, x\} \text{ is a hinge} \}|.$$

The spread of the 2-connected graph G , denoted by δ_G , is defined by

$$\delta_G = \max \{ \delta_v \text{ s.t. } v \in V_G \}.$$

Definition 6.8. A sequence of n hinges $\{x_1, y_1\}, \dots, \{x_n, y_n\}$ is *parallel* if (i) each two hinges are disjoint, and (ii) there is a path $t_1 \dots t_m$ in T such that for each $i \in \{1, \dots, n\}$ there exists $j \in \{1, \dots, m\}$ where $\{x_i, y_i\} \in V_{t_j}$.

For instance, an n -domino where $n \geq 2$ contains $n - 1$ parallel hinges. The following Lemma emphasizes an important aspect of hinges, intuitively it states that hinges *do not cross each other*.

Lemma 6.9. [Theorem IV.22 of [Tut01] or Lemma 8 of [Ric04]]

Let G be a 2-connected graph and let $\{x_1, y_1\}$ be a hinge of G and (U, W) be the 2-separation of G such that $U \cap W = \{x_1, y_1\}$. If $\{x_2, y_2\}$ is an other hinge of G such that $\{x_1, y_1\} \cap \{x_2, y_2\} = \emptyset$, then either $x_2, y_2 \in U$ or $x_2, y_2 \in W$.

Lemma 6.10. Let G be a 2-connected graph and T its Tutte's tree. For all $n \geq 2$, if G contains $4n$ parallel hinges then G has the $(n - 1)$ -domino as a minor.

Proof. Let H be a set of $4n$ hinges of G . First, we construct a graph $\overline{G}$ out of G such that $\overline{G}$ is a minor of G and second, we prove that $\overline{G}$ admits the $(n - 1)$ -domino as a minor.

Lemma 3.3 states that a 2-separator $\{x, y\}$ is a hinge if and only if (i) either there are at least three $[x, y]$ -bridges, or (ii) there are two $[x, y]$ -bridges at least one of them is 2-connected. The construction of $\overline{G}$ follows. For each hinge $\{x, y\} \in H_n$, where there are at least three $[x, y]$ -bridges, do the following operations: (i) delete from G all the $[x, y]$ -bridges which do not contain a hinge in H other than the edge xy ¹⁴, and (ii) add the edge xy to G .

Fact 6.11. The graph $\overline{G}$ constructed above has the following properties:

- (1) $\overline{G}$ is 2-connected,
- (2) $\overline{G}$ is a minor of G ,
- (3) the set H is again a set of hinges of $\overline{G}$, and
- (4) for every $\{x, y\} \in H$, there are exactly two $[x, y]$ -bridges of $\overline{G}$.

Proof. We argue next that the operations (i) and (ii) cited above preserve the properties stated in statements (1),(2),(3), and (4). The statement (1) follows from Lemma 3.5 which states that if $\{x, y\}$ is a 2-separator of G into (U_1, U_2) then $G[U_i] + xy$, for $i = 1, 2$, is also 2-connected. Statement (2) follows from Lemma 3.8 which states that $G[U_i] + xy$ is a minor of G , for $i = 1, 2$. Let us show statement (3). Let $\{x, y\}$ be a hinge in H where $xy \in E_{\overline{G}} \setminus E_G$. On the one hand, $\{x, y\}$ is again a 2-separator in $\overline{G}$. On the other hand, one of the $[x, y]$ -bridges is 2-connected, Lemma 3.5, because it contains the edge xy . Finally, statement (4) holds obviously by construction. This ends the proof of the Fact. $\square$

Second, let us prove that $\overline{G}$ admits the $(n - 1)$ -domino as a minor. Let $H = \{\{a_1, b_1\}, \dots, \{a_{4n}, b_{4n}\}\}$, and v (resp. w) be a vertex in the $[a_1, b_1]$ -bridge (resp. in the $[a_{4n}, b_{4n}]$ -bridge) which does not contain the remaining hinges. Let π_x, π_y be two (simple) disjoint v - w paths in $\overline{G}$. Such paths do exist because $\overline{G}$ is 2-connected, statement (1) of the Fact 6.11. We claim that, for each hinge $\{a_i, b_i\} \in H$, either $a_i \in \pi_x$ and hence $b_i \in \pi_y$ or $a_i \in \pi_y$ and hence $b_i \in \pi_x$. The argument is that each hinge in H is a 2-separator of v from w in $\overline{G}$. Therefore, from now we assume that the set H of hinges is of form

¹⁴Observe that the hinges of H belong to at most two $[x, y]$ -bridges of G .

$H = \{ \{x_i, y_i\}, i = 1, \dots, 4n \}$ such that $x_i \in \pi_x$ and $y_i \in \pi_y$, for $i = 1, \dots, 4n$. Intuitively, the two paths π_x and π_y would play the role the horizontal lines of the domino¹⁵ in question. The main remaining technical part is to construct the vertical lines.

Definition 6.12. Let $i \in \{1, \dots, 4n - 1\}$, and let $G_i \subset \overline{G}$ be the $[x_{i+1}, y_{i+1}]$ -bridge which contains the hinge $\{x_i, y_i\}$ and let $G_{i+1} \subset \overline{G}$ be the $[x_i, y_i]$ -bridge which contains the hinge $\{x_{i+1}, y_{i+1}\}$. We call an $(i, i+1)$ -block the graph $G_i \cap G_{i+1}$. If no confusion will arise, we will call an $(i, i+1)$ -block with simply a block.

Fact 6.13. *If one of G_i and G_{i+1} (given in the previous definition) is 2-connected then the $(i, i+1)$ -block is connected. Moreover, $\overline{G}$ contains at least $2n$ connected blocks.*

Proof. We prove the first statement of the Fact by assuming that G_i is 2-connected. If G_{i+1} is 2-connected then the proof is similar. Let us denote by $\mathcal{B}$ the $(i, i+1)$ -block, and let $v_1, v_2 \in V_{\mathcal{B}}$. We shall show the existence of at least one v_1 - v_2 path in $\mathcal{B}$. Since G_i is 2-connected, then there are at least two disjoint (simple) v_1 - v_2 paths in G_i . Let π_1 and π_2 be such paths. If both π_1 and π_2 are in $\mathcal{B}$ then we have done. Otherwise, assume that π_1 does not belong to $\mathcal{B}$, i.e. π_1 contains a proper subpath which belongs to $G_i \setminus \mathcal{B}$. We claim that π_1 contains both x_i and y_i because $\{x_i, x_{i+1}\}$ is a 2-separation in G_i and the path π_1 is supposed to be simple. We mean that if π_1 visits x_i , and visits some vertices in $G_i \setminus \mathcal{B}$, then it must visit y_i . Therefore, π_1 may be of the form:

$$\pi_1 = v_1 \dots v_p x_i w_1 \dots w_l y_i v_{p+1} \dots v_q v_2,$$

or of the form:

$$\pi_1 = v_1 \dots v_p y_i w_1 \dots w_l x_i v_{p+1} \dots v_q v_2,$$

where $v_j \in V_{\mathcal{B}}$ for $j = 1, \dots, q$ and $w_j \in V_{G_i} \setminus V_{\mathcal{B}}$ for $j = 1, \dots, l$. Since π_1 contains both x_i and y_i , then π_2 contains neither x_i nor y_i , because π_1 and π_2 are supposed to be disjoint. Therefore π_2 belongs to $\mathcal{B}$, i.e. the block $\mathcal{B}$ is connected.

To prove the second statement of the Fact, recall that from the statement (4) of the Fact 6.11 we have the number of $[x_j, y_j]$ -bridges equals 2, for each hinge $\{x_j, y_j\}$ in H . From the statement (ii) of Lemma 3.3 it follows that one of the two $[x_j, y_j]$ -bridges is 2-connected. This implies that one of the $(j-1, j)$ -block and $(j, j+1)$ -block is connected. Hence, $\overline{G}$ does not contain two contiguous¹⁶ blocks which are both not connected. Therefore, $\overline{G}$ contains at least $\frac{4n}{2}$ connected blocks. This ends the proof of the Fact. $\square$

According to the previous Fact, $\overline{G}$ contains at least $2n$ connected blocks. Therefore, $\overline{G}$ contains at least n non contiguous and connected blocks $\mathcal{B}_1, \dots, \mathcal{B}_n$, the latter are pairwise disjoint. Each block $\mathcal{B}_i$, $i = 1, \dots, n$, contains a hinge

¹⁵if the domino is viewed in an horizontal way as depicted in Figure 6.2.

¹⁶Two blocks $(i, i+1)$ and $(j, j+1)$ are contiguous if $i = j+1$ or $j = i+1$.

$\{a_i, b_i\} \in H$, and moreover it contains a a_i - b_i path π_i because it is connected. On the one hand, the graph consisting of the paths $\pi_i, i = 1, \dots, n$, π_x , and π_y is a subgraph of $\overline{G}$. On the other hand, the π_i paths, for $i = 1, \dots, n$, are pairwise disjoint. By contracting each path among π_i , π_x , and π_y in the desired way we get an $(n - 1)$ -domino, see Figure 6.2.




Figure 6.2: The construction of the $(n - 1)$ domino as a minor

The latter has a vertical edge $a_i b_i$ for each hinge $\{a_i, b_i\}$. This ends the proof of Lemma 6.10. $\square$

The following Lemma establishes the relation between the spread, the diameter, and the number of parallel hinges.

Lemma 6.14. *Let G be a 2-connected graph and T be its Tutte's tree. Let δ_G be the spread of G . For all $n \geq 2$, if the diameter of T is greater than $4n \cdot \delta_G$ then G contains at least n parallel hinges.*

Proof. Consider a path in T of length $4n \cdot \delta_G$ of the form

$$t_1 t'_1 \dots t_{2n \cdot \delta_G} t'_{2n \cdot \delta_G} t_{2n \cdot \delta_G + 1}$$

where t_1 is a leaf. One of Tutte's tree properties, see Lemma 3.2, ensures that T enjoys the following properties:

- (a) T is bipartite and moreover every node $t'_i, i = 1, \dots, 2n \cdot \delta_G$ is such $V_{t'_i}$ is a hinge.
- (b) if h_i denotes the hinge $V_{t'_i}$ then
 - (b.1) for all $i = 1, \dots, 2n \cdot \delta_G - 1$, we have either $h_i \cap h_{i+1} = \emptyset$ or $|h_i \cap h_{i+1}| = 1$,
 - (b.2) if $h_p \cap h_q = \emptyset$ where $1 \leq p < q \leq 2n \cdot \delta_G$, then for all $i \leq p$ we have that $h_i \cap h_q = \emptyset$, and
 - (b.3) if $h_p \cap h_q = \emptyset$ where $1 \leq p < q \leq 2n \cdot \delta_G$, then for all $i \geq q$ we have that $h_p \cap h_i = \emptyset$.

Let $H = \{h_1, \dots, 2n\delta_G\}$ be a sequence of hinges where $h_i = V_{t_i}$. Out of H we shall define an undirected graph ∂H in such a way the properties of hinges in H transfer into the properties of ∂H . Define ∂H as follows $V_{\partial H} = \{h^v \mid h \in H\}$, in other words the vertex h^v is just the hinge h viewed as a vertex; and $h_1^v h_2^v \in E_{\partial H}$ if and only if $h_1 \cap h_2 \neq \emptyset$. We state the main properties of the graph ∂H .

Fact 6.15. *Let $\mathcal{B}$ be a 2-connected component of ∂H , and let $m = |V_{\mathcal{B}}|$, then $\mathcal{B}$ is an m -clique.*

Proof. Let $h_1^v, h_2^v \in V_{\mathcal{B}}$. Since $\mathcal{B}$ is 2-connected then it follows by Menger Theorem 2.3 that there exist two disjoint h_1^v - h_2^v paths in $\mathcal{B}$ i.e. there is a cycle in $\mathcal{B}$ containing both h_1^v and h_2^v . Let $h_1^v, h_2^v, \dots, h_m^v h_1^v$ be such a cycle. Assume that $h_1^v h_3^v \notin E_{\mathcal{B}}$, hence $h_1 \cap h_3 = \emptyset$. It follows from property (b.3) above that for all $i \geq 3$ we have $h_1 \cap h_i = \emptyset$. This is a contradiction since already $h_1 \cap h_n \neq \emptyset$ because $h_1^v h_m^v \in E_{\mathcal{B}}$. We deduce that $h_1^v h_3^v \in E_{\mathcal{B}}$. Using similar arguments, we deduce that for all $i = 3, \dots, m-1$ we have $h_1^v h_{m-i}^v \in E_{\mathcal{B}}$. We conclude that for every pair h_1^v, h_2^v of vertices in the component $\mathcal{B}$ there is an edge $h_1^v h_2^v$ in $\mathcal{B}$, implying that $\mathcal{B}$ is an m -clique. This ends the proof of the Fact. $\square$

Fact 6.16. *If ∂H contains an m -clique of vertices $h_1^v, \dots, h_m^v$ then*

$$|\cap_{i=1, \dots, m} h_i| = 1.$$

Proof. The proof is by induction on m .

If $m = 3$, then either $\cap_i h_i \neq \emptyset$ and in this case $|\cap_i h_i| = 1$, or $\cap_i h_i = \emptyset$ and in this case the hinges h_1, h_2, h_3 form a triangle in the following sense: $h_i \cap h_j = \{v_{ij}\}$ for $i, j = 1, 2, 3$ and $i \neq j$. In the latter case, the hinges h_1, h_2, h_3 do not belong to a unique path in T , contradicting the hypothesis that the hinges in question are parallel, condition (ii) of Definition 6.8.

Induction step. Consider a $(m+1)$ -clique of vertices $h_1^v, \dots, h_{m+1}^v$. By induction hypothesis $|\cap_{i=2, \dots, m+1} h_i| = 1$ and $|\cap_{i=1, \dots, m} h_i| = 1$. This implies that $\cap_{i=2, \dots, m+1} h_i = \cap_{i=1, \dots, m} h_i$, and hence $|\cap_{i=1, \dots, m+1} h_i| = 1$. This ends the proof of the Fact. $\square$

Summing up the Facts 6.15 and 6.16 we deduce that the size (i.e. the number of vertices) of the 2-connected components of ∂H is at most δ_G , because, given a 2-connected component $\mathcal{B}$ of vertices $h_1^v, \dots, h_m^v$ the spread of the vertex $\cap_{i=1, \dots, m} h_i$ is m . The following Claim allows us to compute a lower bound of the number of 2-connected components of ∂H .

Claim 6.17. *Let H be a graph. If the size (i.e. number of vertices) of its 2-connected components is at most δ then G has got at least $\lfloor \frac{|V_H|}{\delta} \rfloor$ 2-connected components.*

Proof. Let x be the number of 2-connected components of H . Clearly $|V_H| \leq x\delta$, hence $\frac{|V_H|}{\delta} \leq x$. Therefore H contains at least $\lfloor \frac{|V_H|}{\delta} \rfloor$ 2-connected components. This ends the proof of the Claim. $\square$

Since $|V_{\partial H}| = 2n.\delta_G$ and the size of the 2-connected components of ∂H is at most δ_G , then, according to the Claim, the number of the 2-connected components of ∂H is at least $2n$.

On the one hand, if the two vertices h_1^v and h_2^v belong to two disjoint 2-connected components then the related hinges h_1 and h_2 are disjoint, because otherwise, there is an edge $h_1^v h_2^v$ in ∂H implying that both h_1^v and h_2^v belong to the same 2-connected component, which is a contradiction. On the other hand ∂H contains at least n pairwise disjoint 2-connected components. We conclude that G contains at least n disjoint hinges, the latter belong to the same path in T , and therefore they are parallel. This ends the proof of Lemma 6.14. $\square$

Now we are ready to provide an upper bound of the diameter of Tutte's tree if the related graph has entanglement 3.

Proposition 6.18. *Let G be a 2-connected graph, T be its Tutte's tree, and δ_G be the spread of G . If the entanglement of G is 3 then the diameter of T is at most $2^7.\delta_G$.*

Proof. Let G be as stated in the Proposition with $\mathcal{E}(G) = 3$. If the diameter of G is $c.\delta_G$ then it follows from the Lemma 6.14 that G contains at least $\lfloor c \setminus 4 \rfloor$ parallel hinges. Therefore, from Lemma 6.10 we get that G contains a c' -domino $D_{c'}$ as a minor, where $c' = \lfloor \lfloor c \setminus 4 \rfloor \setminus 4 \rfloor - 1$. Hence $\mathcal{E}(D_{c'}) \leq \mathcal{E}(G)$ by Theorem 4.3 which states that the class of undirected graphs of entanglement at most k is minor ideal. From Lemma 6.5, a 14-domino has entanglement strictly greater than 3. We conclude that $c' \leq 14$ and hence $c \leq 2^7$. $\square$

Now we are ready to state the main result of this paper. To this goal, we define the *interface* of a torso τ_t as

$$I_t = \{ v \in V_t \mid \exists tt' \in E_T \text{ s.t } v \in V_{t'} \}.$$

Theorem 6.19. *Let G be a 2-connected graph and let $(T, (V_t)_{t \in T})$ be the Tutte decomposition of G . If the entanglement of G is 3 then for every 3-connected torso τ_t of G the following hold:*

1. *there exists $B_3 \subseteq V_t$ where $|B_3| = 3$ such that (τ_t, B_3) is an abstract 3-molecule,*
2. *$I_t \subseteq B_3$, and*
3. *if δ_G is the spread of G , then the diameter of T is at most $2^7.\delta_G$.*

Proof. Let G be as stated and τ_t a 3-connected torso of G .

1. On the one hand, Lemma 3.7 states that τ_t is a minor of G . It follows from Theorem 4.3 that $\mathcal{E}(\tau_t) \leq 3$. On the other hand, since τ_t is 3-connected then we get from Lemma 4.4 that $3 \leq \mathcal{E}(\tau_t)$, therefore $\mathcal{E}(\tau_t) = 3$. Hence, from Lemma 5.7 we deduce that there exists $B_3 \subset V_{\tau_t}$ with $|B_3| = 3$ such that (τ_t, B_3) is an abstract 3-molecule.

2. If the 3-molecule τ_t is not ambiguous, then the property $I_t \subseteq B_3$ follows from statement (ii) of Corollary 6.3. If τ_t is not ambiguous, then it suffices to generalize Lemma 6.1 as follows:

Lemma 6.20. *Let ϑ be a 3-molecule, and let each C_3, C'_3, C_3 and C''_3 be a 3-cycle. Let*

$$G = ((\vartheta +_{v_1 w_1} C_3) +_{v_2 w_2} C'_3) +_{v_3 w_3} C''_3,$$

where $+$ is the 2-Sum operator and $v_i w_i \in E_{\vartheta}$, $i = 1, 2, 3$, $v_1 w_1 \in E_{C_3}$, $v_2 w_2 \in E_{C'_3}$, and $v_3 w_3 \in E_{C''_3}$. If the entanglement of G is again 3 then this implies that the vertices $v_i, w_i, i = 1, 2, 3$ belong to a minimal edge cover set of ϑ .

3. The condition on the diameter of T follows from Proposition 6.18.

□

7 Towards sufficient conditions on Tutte's tree

Theorem 6.19 provides some necessary conditions on Tutte's tree to be a tree decomposition of a 2-connected graph of entanglement 3. However, these conditions are not sufficient, since the 14-domino is a counter example. Technically speaking, the 14-domino is not a worth counter example because the computation of the exact value of Tutte's tree would allow us to obtain sufficient conditions. To be more precise, the exact value of the diameter would essentially depend on, besides the spread, the length of each cycle which constitutes a torso of Tutte's tree. In the sequel we shall tell something about the entanglement of 2-connected graphs for which Tutte's tree satisfies conditions (1) and (2) of Theorem 6.5 and the hinges of G are organized into a path-like structure in the following sense: there exists a path $t_1, \dots, t_n$ in T such that for each hinge h of G there exists t_i where $h = V_{t_i}$, $i \in \{1, \dots, n\}$.

Proposition 7.1. *Let G be a 2-connected graph and T be its Tutte's tree. If T satisfies conditions (1) and (2) of Theorem 6.5 and if the hinges of G are organized into a path-like structure, then G has entanglement at most 4.*

Proof. The proof of This Proposition is essentially an adaptation of the proof of Proposition 8.8.1 of [Bel08], the latter states that the entanglement of the n -domino is at most 4.

Let G and T be as stated in the Proposition. We shall describe a winning strategy for Cops in $\mathcal{E}(G, 4)$.

Let $H = \{h_i \mid i = 0, n\}$ be the sequence of hinges of G . Cops strategy in $\mathcal{E}(G, 4)$ is described by means of the following steps:

- Step 1. By playing on the hole graph and using just 2 cops, occupy a hinge of the form h_j where $0 \leq j \leq n$. At this moment, Thief goes either to an

h_j -bridge which does not contain any hinge, or he goes to an h_j -bridge which contain at least one hinge. Observe that there exist at most two h_j -bridges which contain hinges because the bridges of G are organized into a path-like structure. In the first case Cops use the third cop to catch Thief in the h_j -bridge, the latter is either a path, a cycle, or a 3-molecule by assumptions 1 and 2 of Theorem 6.19. In the second case, iterate the strategy from Step 2.

- Step 2. By playing on the related h_j -bridge – where 2 cops are placed on h_j – Cops use the other 2 cops to occupy a hinge of form h_i . At this moment, Thief has 3 possibilities: (1) he goes to one of the h_i -bridges which does not contain any hinge, and therefore he will be trapped, (2) he goes to the h_i -bridge which does not contain the hinge h_j , and in this case iterate the strategy from Step 2 (up to symmetry) by keeping the two cops on h_i and using the two cops on h_j , or (3) he goes to the h_i -bridge which contains the hinge h_j , and in this case iterate the strategy from Step 3.
- Step 3. By playing on the h_i -bridge $\cap$ h_j -bridge which contains both h_i and h_j – where the 4 cops are placed on $h_i \cup h_j$ ¹⁷ and Thief is on vertex $x_i \in h_i$, (the other positions of Thief are handled by symmetry) – Cops use the two cops on h_k to occupy a hinge of the form h_p where $i < p < j$ and k equals j if $j \leq n - i$ and equals i otherwise. Besides, Cops do not remove the 2 cops on $h_{\bar{k}}$, where $\bar{k} = i$ if $k = j$ and $\bar{k} = j$ if $k = i$. At this moment, i.e. from the position $x_p \in h_p$, Thief has three possibilities: either (1) he goes to an h_p -bridge which does not contain any hinge, and therefore he will be trapped, (2) he goes to the h_p -bridge which contains hinges but not the $h_{\bar{k}}$ one, and in this case iterate the strategy from Step 2, or (3) he goes to the h_p -bridge $\cap$ $h_{\bar{k}}$ -bridge and in this case iterate the strategy from Step 3.

It remains to argue that (i) Cops strategy described in each step is realizable and (ii) this strategy would not be iterated infinitely often. To prove statement (i), it is enough to prove the following Fact.

Fact 7.2. *Let G be a 2-connected graph and T be its Tutte's tree. If T satisfies conditions (1) and (2) of Theorem 6.19 and if the hinges of G are organized into a path-like structure, then in the game $\mathcal{E}(G, 2)$ Cops have a strategy to occupy a hinge of G or they win.*

Proof. Let $H = \{ \{ x_i, y_i \}, i = 0, n \}$ be the sequence of hinges of G .

First, if Thief restricts his moves on some torso $\tau_t, t \in V_t$, then we distinguish two cases according to the nature of τ_t :

Case (i). if τ_t is a cycle of the form $v_1 v_2 \dots v_m v_1$, and Thief is walking from v_1 to v_m , then Cops strategy consists in putting the first cop on v_1 and following Thief with the second cop, until a vertex of some hinge $h_i \subset \{ v_1, \dots, v_m \}$ is

¹⁷We can assume w.l.g that $h_i \cap h_j = \emptyset$, otherwise Thief will be easily trapped.

occupied by a cop, and then Cops do not move this cop and follow Thief with the other cop until the second vertex of the hinge h_i is occupied by the second cop, and we have done.

Case (ii). if τ_t is 3- connected then it is a 3-molecule by condition (1) of Theorem 6.19 and the interface I_t of τ_t belongs to a minimal edges cover set of τ_t by condition (2) of the same Theorem. In this case, Cops strategy consists in skipping on the vertices $V_{\tau_t} \setminus I_t$ and using the two cops to occupy two vertices in I_t . These two vertices obviously constitute a hinge of G , and we have done.

Second, if Thief does not restrict his moves on the same torso, i.e. he moves on a sequence of torsos $\tau_{t_1}, \tau_{t_2}, \dots$ where $t_1, t_2, \dots \in V_T$, then Cops strategy consists in placing the first cop on the vertex of $\tau_{t_1} \cap \tau_{t_2}$ which is visited by Thief (recall that $\tau_{t_1} \cap \tau_{t_2}$ is a hinge), placing the second cop on the vertex – visited by Thief – of $\tau_{t_2} \cap \tau_{t_3}$, replacing the first cop to the vertex – visited by Thief – of $\tau_{t_3} \cap \tau_{t_4}$, ... until some torso τ_{t_i} is visited twice by Thief. Observe that there are two cops placed on $\tau_{t_i} \cap \tau_{t_{i+1}}$. Since $\tau_{t_i} \cap \tau_{t_{i+1}}$ is a hinge of G , we have done. This completes the proof of the Fact. $\square$

To show statement (ii) above it is enough to observe that whenever a step is revisited a second time, then the value $|i - j|$ is strictly lower than that of the first time. This ends the proof of Proposition 7.1. $\square$

As a direct consequence of the previous Proposition, the n -domino has entanglement at most 4 and therefore it follows from Lemma 6.5 that the entanglement of the n -domino, where $n \geq 14$, is exactly 4.

Acknowledgement. We acknowledge Luigi Santocanale for helpful discussions on the topic.

References

- [Bel08] Walid Belkhir. *Algebra and Combinatorics of Parity Games*. PhD thesis, Université de Provence, Dec 2008. <http://www.lif.univ-mrs.fr/~belkhir/these.pdf>.
- [BG05] Dietmar Berwanger and Erich Grädel. Entanglement—a measure for the complexity of directed graphs with applications to logic and games. In *LPAR 2005*, volume 3452 of *LNCS*, pages 209–223. Springer, 2005.
- [BGL07] Dietmar Berwanger, Erich Grädel, and Giacomo Lenzi. The variable hierarchy of the μ -calculus is strict. *Theory Comput. Syst.*, 40(4):437–466, 2007.
- [BJG01] Jørgen Bang-Jensen and Gregory Gutin. *Digraphs*. Springer Monographs in Mathematics. Springer-Verlag London Ltd., London, 2001. Theory, algorithms and applications.

- [BS07] Walid Belkhir and Luigi Santocanale. Undirected graphs of entanglement 2. In V. Arvind and S. Prasad, editors, *FSTTCS 2007*, volume 4855 of *Lect. Not. Comp. Sci.*, pages 508–519. Springer, 2007.
- [BS08] Walid Belkhir and Luigi Santocanale. The variable hierarchy for the lattice μ -calculus. In Ilario Cervesato, Helmut Veith, and Andrei Voronkov, editors, *LPAR 2008*, *Lect. Not. Comp. Sci.*, pages 605–620, 2008.
- [Die05] Reinhard Diestel. *Graph theory*, volume 173 of *Graduate Texts in Mathematics*. Springer-Verlag, Berlin, third edition, 2005.
- [DSS95] Carl Droms, Brigitte Servatius, and Herman Servatius. The structure of locally finite two-connected graphs. *Electron. J. Combin.*, 2:Research Paper 17, approx. 10 pp. (electronic), 1995.
- [FPR99] Paola Festa, Panos M. Pardalos, and Mauricio G. C. Resende. Feedback set problems. In *Handbook of combinatorial optimization, Supplement Vol. A*, pages 209–258. Kluwer Acad. Publ., Dordrecht, 1999.
- [Hoh92] Walter Hohberg. The decomposition of graphs into k -connected components. *Discrete Math.*, 109(1-3):133–145, 1992.
- [Koz83] Dexter Kozen. Results on the propositional μ -calculus. *Theoret. Comput. Sci.*, 27(3):333–354, 1983.
- [Ric04] R. Bruce Richter. Decomposing infinite 2-connected graphs into 3-connected components. *Electr. J. Comb.*, 11(1), 2004.
- [RS86] Neil Robertson and P. D. Seymour. Graph minors. II. Algorithmic aspects of tree-width. *J. Algorithms*, 7(3):309–322, 1986.
- [RSŠ94] R. B. Richter, P. D. Seymour, and J. Širáň. Circular embeddings of planar graphs in nonspherical surfaces. *Discrete Math.*, 126(1-3):273–280, 1994.
- [Sey81] P. D. Seymour. Some applications of matroid decomposition. In *Algebraic methods in graph theory, Vol. I, II (Szeged, 1978)*, volume 25 of *Colloq. Math. Soc. János Bolyai*, pages 713–726. North-Holland, Amsterdam, 1981.
- [Tut66] W. T. Tutte. *Connectivity in graphs*. Mathematical Expositions, No. 15. University of Toronto Press, Toronto, Ont., 1966.
- [Tut01] W. T. Tutte. *Graph theory*, volume 21 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2001. With a foreword by Crispin St. J. A. Nash-Williams, Reprint of the 1984 original.
- [Wag37] K. Wagner. Über eine Eigenschaft der ebenen Komplexe. *Math. Ann.*, 114(1):570–590, 1937.